


Redox reactions – Galvanic cells

Lesson 4a

Galvanic cells convert chemical potential energy into electrical energy. A redox reaction takes place where the oxidation and reduction half reactions are separated and electrons forced to travel from the oxidation half reaction to the reduction half reaction via an external circuit.




A few things to note when referring to galvanic cells.

- Energy is converted from chemical to electrical
- The oxidant and reductant are separated into two half-cells
- Oxidation occurs at the electrode known as the anode
- The anode has a negative polarity
- Reduction occurs at the electrode known as the cathode
- The cathode has a positive electrode
- Electrons travel spontaneously from anode to cathode ie. from negative to positive via an external circuit
- The two half-cells are connected by a salt bridge containing an ionic substance, usually $\text{KNO}_3(\text{aq})$. Positive ions (K^+) flow from the salt bridge into the half cells containing the cathode while negative ions (NO_3^-) flow from the salt bridge into the half-cell containing the anode.
- In order for a spontaneous reaction to occur the oxidant must be higher than the reductant on the E° table, as shown on the right.
It is important to note, that having the oxidant higher than reductant indicates that a reaction will occur, however, the rate of the reaction may be so slow that no reaction is observed.
- The half-cell containing the anode has the reductant and its conjugate oxidant, in this case Zn/Zn^{2+} , and the half-cell containing the cathode has the oxidant and its conjugate reductant, in this case Cu^{2+}/Cu
- If one of the species in the half-cell is a metal it is often used as the electrode. In this case copper and zinc are both used as electrodes. However a conducting material, such as graphite, can also be used.
- The voltage produced by the galvanic cell, at standard conditions (25°C , 1 atm pressure and 1.0 M concentration) is given by the formula below
 Cell potential difference = higher half-cell E° – lower half-cell E°
 In this case it is cell potential = $+0.34\text{V} - -0.76\text{V} = 1.10\text{V}$



oxidant	$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$	+0.34
	$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}(\text{aq})$	+0.15
	$\text{S}(\text{s}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$	+0.14
	$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	0.00
	$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Pb}(\text{s})$	-0.13
	$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}(\text{s})$	-0.14
	$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ni}(\text{s})$	-0.23
	$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Co}(\text{s})$	-0.28
	$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Fe}(\text{s})$	-0.44
	$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Zn}(\text{s})$ reductant	-0.76

- 1) Consider the galvanic cells shown on the right.
For each cell give the cell potential and label the:
- Direction of electron flow
 - Anode and cathode
 - Direction of K^+ ions flowing from the salt bridge
 - Direction of NO_3^- ions flowing from the salt bridge



- 2) Predict if a spontaneous reaction will occur in the galvanic cells shown on the right


No

$Ag^+(aq) + e^- \rightleftharpoons Ag(s)$	Ag is the reductant no oxidant above the reductant	+0.80
$Fe^{3+}(aq) + e^- \rightleftharpoons Fe^{2+}(aq)$		+0.77
$O_2(g) + 2H^+(aq) + 2e^- \rightleftharpoons H_2O_2(aq)$		+0.68
$I_2(s) + 2e^- \rightleftharpoons 2I^-(aq)$		+0.54
$O_2(g) + 2H_2O(l) + 4e^- \rightleftharpoons 4OH^-(aq)$		+0.40
$Cu^{2+}(aq) + 2e^- \rightleftharpoons Cu(s)$		+0.34
$Sn^{4+}(aq) + 2e^- \rightleftharpoons Sn^{2+}(aq)$	Sn ⁴⁺ is the only oxidant and it is below the reductant	+0.15

3) The overall redox reaction in a galvanic cell is given below.
 $\text{Cu(s)} + \text{Cl}_2(\text{g}) \rightarrow \text{Cu}^{2+}(\text{aq}) + 2\text{Cl}^-(\text{aq})$

- i. Give the oxidation and reduction half equations
- ii. Draw a fully labelled diagram, as in 1) above, of the galvanic cell.

(Consider the shape of a gas electrode)



4) Consider the Galvanic cell shown on the right.
 What type of energy is produced by this device if at all.

Ag⁺ is the oxidant that is placed in the same half-cell as the reductant Cu.

A reaction will occur as the oxidant is higher than the reductant, as shown on the right.

Since the reaction takes place in the same half-cell electrons are exchanged directly and are not forced to travel through an external circuit. Heat energy is produced and not electrical energy.

$\text{Ag}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s})$
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$
$\text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2(\text{aq})$
$\text{I}_2(\text{s}) + 2\text{e}^- \rightleftharpoons 2\text{I}^-(\text{aq})$
$\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightleftharpoons 4\text{OH}^-(\text{aq})$
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$